

Contents lists available at ScienceDirect

Journal of Algebra





Research Paper

Twisted bimodules and universal enveloping algebras associated to VOAs



Jianzhi Han^a, Yukun Xiao^{b,c,*}, Shun Xu^d

- ^a School of Mathematical Sciences, Key Laboratory of Intelligent Computing and Applications (Ministry of Education), Tongji University, Shanghai, 200092, China
 ^b School of Mathematics and Statistics, Qingdao University, Qingdao, 266071, China
- ^c Qingdao International Mathematics Center, Qingdao, 266071, China
- d School of Mathematical Sciences, Tongji University, Shanghai, 200092, China

ARTICLE INFO

Article history: Received 23 June 2024 Available online 30 October 2024 Communicated by Alberto Elduque

MSC: 17B69

Keywords: Vertex operator algebras Associative algebras Bimodules Universal enveloping algebras

ABSTRACT

For any vertex operator algebra V, finite automorphism g of V of order T and $m, n \in (1/T)\mathbb{Z}_+$, we construct a family of associative algebras $A_{g,n}(V)$ and $A_{g,n}(V)-A_{g,m}(V)$ -bimodules $A_{g,n,m}(V)$ from the point of view of representation theory. We prove that the algebra $A_{g,n}(V)$ is identical to the algebra $A_{g,n}(V)$ constructed by Dong, Li and Mason, and that the bimodule $A_{g,n,m}(V)$ is identical to $A_{g,n,m}(V)$ which was constructed by Dong and Jiang. We also prove that the $A_{g,n}(V)-A_{g,m}(V)$ -bimodule $A_{g,n,m}(V)$ is isomorphic to $U(V[g])_{n-m}/U(V[g])_{n-m}^{m-1/T}$, where $U(V[g])_k$ is the subspace of degree k of the $(1/T)\mathbb{Z}$ -graded universal enveloping algebra U(V[g]) of V with respect to g and $U(V[g])_k^l$ is some subspace of $U(V[g])_k$. And we show that all these bimodules $A_{g,n,m}(V)$ can be defined in a simpler way.

© 2024 Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

^{*} Corresponding author.

E-mail addresses: jzhan@tongji.edu.cn (J. Han), ykxiao@qdu.edu.cn (Y. Xiao), shunxu@tongji.edu.cn (S. Xu).

1. Introduction

The representation theory of vertex operator algebras is quiet different from that of classical algebras because of the appearance of twisted modules. Among all representations of a vertex operator algebra, admissible twisted modules are the most important ones. Recall that for a vertex operator algebra V and an automorphism g of V of finite order T, an admissible g-twisted V-module M is $(1/T)\mathbb{Z}_+$ -graded: $M = \bigoplus_{i \in (1/T)\mathbb{Z}_+} M_i$ (cf. [7]). Thus, in order to study M it is vital to determine all $\operatorname{Hom}(M_i, M_j)$ for $i, j \in (1/T)\mathbb{Z}_+$. In fact, a series of associative algebras $A_{g,n}(V)$ was introduced (see [7,8]) for which there is an algebra homomorphism from $A_{g,n}(V)$ to $\operatorname{Hom}(M_i, M_i)$ for any $i \leq n$. And in generally, to study these $\operatorname{Hom}(M_i, M_j)$, a series of $A_{g,n}(V) - A_{g,m}(V)$ -bimodules $A_{g,n,m}(V)$ for $m, n \in (1/T)\mathbb{Z}_+$ was constructed by Dong and Jiang in [4], for which there is an $A_{g,n}(V) - A_{g,m}(V)$ -bimodule homomorphism from $A_{g,n,m}(V)$ to $\operatorname{Hom}(M_{m-l}, M_{n-l})$ for any $0 \leq l \leq \min\{m, n\}$. Thus, in this sense bimodules $A_{g,n,m}(V)$ are generalizations of these associative algebras $A_{g,n}(V)$ (see [4]).

There are several kinds of associative algebras associated to vertex operator (super)algebras (see [2,6,9,14,18,20,22,23,28]); for the twisted case one can refer to [10,11,19,24,25].

From the construction, the bimodule $A_{g,n,m}(V)$ is the quotient of V by $O_{g,n,m}(V)$. Thus, for better understanding how these $A_{g,n,m}(V)$ can be used to study admissible twisted modules, a key step is to study the subspace $O_{g,n,m}(V)$. Intuitively, $O_{g,n,m}(V)$ should be closely related to twisted modules. This is indeed the case when g=1. It was proved in [15] that $O_{1,n,m}(V)$ can also be defined from representations of V. This drives us to do so for general automorphisms. In fact we shall consider another subspace $\mathcal{O}_{g,n,m}(V)$ from the perspective of representation theory and define $\mathcal{A}_{g,n,m}(V)$ as the quotient of V by $\mathcal{O}_{g,n,m}(V)$. In this way, we obtain a series of associative algebras $\mathcal{A}_{g,n}(V)$. As a result, $\mathcal{A}_{g,n,m}(V)$ becomes an $\mathcal{A}_{g,n}(V)-\mathcal{A}_{g,m}(V)$ -bimodule (see Theorem 3.5). And for any $\mathcal{A}_{g,m}(V)$ -module U, we use a different way from [15] to show that $\bigoplus_{n\in(1/T)\mathbb{Z}_+}\mathcal{A}_{g,n,m}(V)\otimes_{\mathcal{A}_{g,m}(V)}U$ is an admissible g-twisted modules with some universal property (see Theorem 3.7). Based on this universal property we can show that $\mathcal{A}_{g,n,m}(V)$ is identical to $A_{g,n,m}(V)$ (see Theorem 5.2). Thus, all bimodules can be reconstructed from the perspective of representation theory. A generalization of twisted bimodules is constructed in [27] also from this perspective.

It is well known that the most powerful tool in the representation theory of Lie algebras is the universal enveloping algebra. As for a vertex operator algebra V and an automorphism g, a weak version of such universal enveloping algebra U(V[g]) also exists: every weak g-twisted V-module is automatically a U(V[g])-module. In fact, these universal enveloping algebras have close connection with $A_{g,n}(V)$ and bimodules $A_{g,n,m}(V)$. Frenkel and Zhu [14] pointed out that Zhu's algebra can be identified with some quotient of U(V[1]). It was proved in [17] that all $A_{g,n}(V)$ for $n \in \mathbb{Z}_+$ are some quotients of the universal enveloping algebra U(V[g]) for the case g = 1 and in [16] for the general finite automorphism g. Furthermore, the bimodule $A_{1,n,m}(V)$ was also proved to be some quotient of U(V[1]) (see [15]). In this present paper, we are going to consider the general g

and show that the $A_{g,n}(V)-A_{g,m}(V)$ -bimodule $A_{g,n,m}(V)$ is some quotient of U(V[g]) (see Theorem 5.2).

As mentioned above, the bimodule $A_{g,n,m}(V)$ is the quotient of V by $O_{g,n,m}(V)$. For some technical reason, $O_{g,n,m}(V)$ is defined as the sum of three subspaces $O'_{g,n,m}(V)$, $O''_{g,n,m}(V)$ and $O'''_{g,n,m}(V)$. But it was conjectured that $O_{g,n,m}(V) = O'_{g,n,m}(V)$ (see [3]), that is, $O''_{g,n,m}(V)$ and $O'''_{g,n,m}(V)$ are superfluous. In this present paper we shall approach to this conjecture. More precisely, we shall show that $O''_{g,n,m}(V)$ is superfluous and $O'_{g,n,m}(V)$ can be replaced by its subspace $\bigoplus_{s \neq \bar{m} - \bar{n} \mod T} V^s + L_{n,m}(V)$, or equivalently,

$$O_{g,n,m}(V) = \bigoplus_{s \not\equiv \bar{m} - \bar{n} \bmod T} V^s + L_{n,m}(V) + O''_{g,n,m}(V) \quad \text{(see Theorem 6.8)}.$$

In this way, we can refine all these bimodules $A_{q,n,m}(V)$.

The paper is organized as follows: In Section 2, we recall the definition of $A_{g,n}(V)-A_{g,m}(V)$ -bimodule $A_{g,n,m}(V)$ and the construction of the universal admissible g-twisted V-module M(U). In Section 3, we construct the associative algebras $A_{g,n}(V)$, the $A_{g,n}(V)-A_{g,m}(V)$ -bimodules $A_{g,n,m}(V)$ and use these bimodules $A_{g,n,m}(V)$ to construct admissible g-twisted V-modules M(U) which have some universal property. In Section 4, we first recall the definition of universal enveloping algebra U(V[g]) of a vertex operator algebra V with respect to a finite automorphism g. Then we obtain another $A_{g,n}(V)-A_{g,m}(V)$ -bimodule $U(V[g])_{n-m}/U(V[g])_{n-m}^{m-1/T}$ and construct universal admissible g-twisted V-modules M(U). In Section 5, we prove that $A_{g,n}(V)$ and $A_{g,n}(V)$ are identical as associative algebras; $A_{g,n,m}(V)$, $A_{g,n,m}(V)$ and $U(V[g])_{n-m}/U(V[g])_{n-m}^{m-1/T}$ are isomorphic to each other as $A_{g,n}(V)-A_{g,m}(V)$ -bimodules. The Section 6 is devoted to refining the definition of the bimodules $A_{g,n,m}(V)$.

We assume that the reader is familiar with the basic knowledge on the vertex operator algebra theory such as the definition of vertex operator algebra (cf. [1], [5], [12], [13], [21]) and the definitions of weak and admissible twisted modules (cf. [7,8]).

2. $A_{g,n}(V) - A_{g,m}(V)$ -bimodule $A_{g,n,m}(V)$

Let $V=\oplus_{n\in\mathbb{Z}}V_n$ be a vertex operator algebra and g an automorphism of V of finite order T. Then we have the decomposition $V=\bigoplus_{r=0}^{T-1}V^r$, where $V^r=\{v\in V\mid gv=e^{-2\pi ri/T}v\}$ (here and only here i represents the imaginary unit). For any n, elements u in V_n are called homogenous and we define wt u=n. So when wt u appears we always assume that u is homogenous.

Let M be a weak g-twisted V-module. Recall from [7] (see also [13]) that for $u \in V^r$, $v \in V$ and $w \in M$, the twisted Jacobi identity

$$\begin{split} z_{0}^{-1}\delta\left(\frac{z_{1}-z_{2}}{z_{0}}\right)Y_{M}\left(u,z_{1}\right)Y_{M}\left(v,z_{2}\right)-z_{0}^{-1}\delta\left(\frac{z_{2}-z_{1}}{-z_{0}}\right)Y_{M}\left(v,z_{2}\right)Y_{M}\left(u,z_{1}\right)\\ &=z_{2}^{-1}\left(\frac{z_{1}-z_{0}}{z_{2}}\right)^{-r/T}\delta\left(\frac{z_{1}-z_{0}}{z_{2}}\right)Y_{M}\left(Y\left(u,z_{0}\right)v,z_{2}\right) \end{split} \tag{2.1}$$

is equivalent to the weak associativity

$$(z_0 + z_2)^{l + \frac{r}{T}} Y_M (u, z_0 + z_2) Y_M (v, z_2) w = (z_2 + z_0)^{l + \frac{r}{T}} Y_M (Y (u, z_0) v, z_2) w$$
 (2.2)

where l is a nonnegative integer, and the commutator formula

$$[Y_M(u, z_1), Y_M(v, z_2)] = \operatorname{Res}_{z_0} z_2^{-1} \left(\frac{z_1 - z_0}{z_2}\right)^{-r/T} \delta\left(\frac{z_1 - z_0}{z_2}\right) Y_M(Y(u, z_0) v, z_2).$$
(2.3)

In fact, the twisted Jacobi identity can be replaced directly by the weak associativity (cf. [21,26]). The following theorem is stated in [26], the detailed proof of which is left in the appendix of the present paper.

Theorem 2.1. [26] Let $(V, Y, \mathbf{1})$ be a vertex algebra and T a positive integer. Let M be a vector space and let $Y_M(\cdot, x)$ be a linear map from V to $(\operatorname{End} M)[[x^{\frac{1}{T}}, x^{-\frac{1}{T}}]]$ such that $Y_M(\mathbf{1}, x) = \operatorname{id}_M$ and $Y_M(v, x)w \in M((x^{\frac{1}{T}}))$ for $v \in V$ and $w \in M$. Set

$$V^r = \left\{ v \in V \mid Y_M(v, x)w \in x^{-\frac{r}{T}}M((x)) \text{ for any } w \in M \right\}$$

for $r \in \mathbb{Z}$. Then the twisted Jacobi identity (2.1) for $u \in V^r, v \in V$ and $w \in M$ is equivalent to the weak associativity (2.2).

For any $n \in (1/T)\mathbb{Z}$, define a linear map $o_n(\cdot): V \to \operatorname{End} M$ sending each element $v \in V$ to $v_{\operatorname{wt} v - 1 + n}$ and by [8] also define

$$\begin{split} \Omega_n(M) &= \{ w \in M \mid v_{\text{wt} \, v - 1 + i} w = 0 \text{ for all } v \in V \text{ and } n < i \in (1/T)\mathbb{Z} \} \\ &= \{ w \in M \mid o_{n+i}(v)w = 0 \text{ for all } v \in V \text{ and } 0 < i \in (1/T)\mathbb{Z} \} \,. \end{split}$$

For short, write $o_n(\cdot)$ as $o(\cdot)$ if n=0. And in this paper, for any two formal variables x, y and any $\alpha \in \mathbb{R}$, we define

$$(x+y)^{\alpha} = \sum_{i \in \mathbb{Z}_+} \frac{\alpha(\alpha-1)\cdots(\alpha-i+1)}{i!} y^i x^{\alpha-i}.$$

Then one can see that

$$(x+y)^{\alpha}(x+y)^{\beta} = (x+y)^{\alpha+\beta}$$
 for any $\alpha, \beta \in \mathbb{R}$.

Proposition 2.2. Let W be a weak g-twisted V-module, $u \in V^r, v \in V^s, p \in r/T + \mathbb{Z}, q \in s/T + \mathbb{Z}$ and $w \in W$. Let $l \in \mathbb{Z}_+$ be such that

$$u_n w = 0$$
 for $n > l + r/T$.

and $k \in \mathbb{Z}_+$ such that

$$v_n w = 0$$
 for $n > k + q$.

And also let $t \in \mathbb{Z}$ be such that

$$u_n v = 0$$
 for $n > t$.

Then

$$u_p(v_q w) = \sum_{i=0}^k \sum_{j=0}^N \binom{p-l-r/T}{i} \binom{l+r/T}{j} (u_{p-l-r/T-i+j} v)_{q+l+r/T+i-j} w,$$

where $N = \max\{t - p + l + r/T + k, 0\}.$

Proof. By (2.2), $z_1^p z_2^q Y(u, z_1) Y(v, z_2) w \in W[[z_1, z_1^{-1}, z_2, z_2^{-1}]]$ and following the proof of [21, Proposition 4.5.7] we see that

$$\begin{split} &u_p(v_q w) \\ &= \operatorname{Res}_{z_1} \operatorname{Res}_{z_2} z_1^p z_2^q Y(u,z_1) Y(v,z_2) w \\ &= \operatorname{Res}_{z_0} \operatorname{Res}_{z_2} (z_0 + z_2)^p z_2^q Y(u,z_0 + z_2) Y(v,z_2) w \\ &= \operatorname{Res}_{z_0} \operatorname{Res}_{z_2} (z_0 + z_2)^{p-l-r/T} z_2^q \big((z_0 + z_2)^{l+r/T} Y(u,z_0 + z_2) Y(v,z_2) w \big) \\ &= \operatorname{Res}_{z_0} \operatorname{Res}_{z_2} (z_0 + z_2)^{p-l-r/T} z_2^q \big((z_2 + z_0)^{l+r/T} Y(Y(u,z_0)v,z_2) w \big) \\ &= \operatorname{Res}_{z_0} \operatorname{Res}_{z_2} \sum_{i=0}^k \binom{p-l-r/T}{i} z_0^{p-l-r/T-i} z_2^{i+q} \big((z_2 + z_0)^{l+r/T} Y(Y(u,z_0)v,z_2) w \big), \end{split}$$

which immediately gives the desired formula.

As for these $\Omega_m(M)$ we have:

Lemma 2.3.
$$o_n(v)\Omega_m(M) \subset \Omega_{m-n}(M)$$
 for $m, n \in (1/T)\mathbb{Z}$ and $v \in V$.

Proof. Without loss of generality we may assume that $v \in V^s$. Take any $u \in V^r$. Then for $0 < i \in (1/T)\mathbb{Z}$ and $w \in \Omega_m(M)$, according to Proposition 2.2, we can choose proper $k, l, N \in \mathbb{Z}_+$ and obtain

$$o_{m-n+i}(u)o_{n}(v)w = u_{\text{wt } u+m-n+i-1}(v_{\text{wt } v+n-1}w)$$

$$= \sum_{p=0}^{k} \sum_{q=0}^{N} {\text{wt } u+m-n+i-1-l-r/T} \binom{l+r/T}{q}$$

$$\times (u_{\text{wt } u+m-n+i-1-l-r/T-p+q}v)_{\text{wt } v+n-1+l+r/T+p-q}w$$

$$= \sum_{p=0}^{k} \sum_{q=0}^{N} \left(wt \, u + m - n + i - l - r/T \right) \binom{l+r/T}{q} o_{m+i} (u_{wt \, u+t-n+i-1-l-r/T-p+q} v) w$$

$$= 0.$$

proving the lemma. \Box

For any $n \in (1/T)\mathbb{Z}_+$, there exists an $\bar{n} \in \{0, 1, \dots, T-1\}$ such that $n = \lfloor n \rfloor + \bar{n}/T$, where $\lfloor \cdot \rfloor$ is the floor function. For $0 \leq r \leq T-1$, define $\delta_i(r) = 1$ if $r \leq i \leq T-1$ and $\delta_i(r) = 0$ if i < r; and set $\delta_i(T) = 0$.

Now for $u \in V^r, v \in V$ and $m, n, p \in (1/T)\mathbb{Z}$, define the product $*_{g,m,p}^n$ on V as follows:

$$u *_{g,m,p}^{n} v = \sum_{i=0}^{\lfloor p \rfloor} (-1)^{i} \begin{pmatrix} \lfloor m \rfloor + \lfloor n \rfloor - \lfloor p \rfloor - 1 + \delta_{\bar{m}}(r) + \delta_{\bar{n}}(T-r) + i \\ i \end{pmatrix}$$
$$\cdot \operatorname{Res}_{z} \frac{(1+z)^{\operatorname{wt} u - 1 + \lfloor m \rfloor + \delta_{\bar{m}}(r) + r/T}}{z^{\lfloor m \rfloor + \lfloor n \rfloor - \lfloor p \rfloor + \delta_{\bar{m}}(r) + \delta_{\bar{n}}(T-r) + i}} Y(u, z) v,$$

if $m, n, p \in (1/T)\mathbb{Z}_+$ and $\bar{p} - \bar{n} \equiv r \mod T$; and $u *_{g,m,p}^n v = 0$ otherwise. Denote $*_{g,m,p}^n$ by $\bar{*}_{g,m}^n$ if p = n and by $*_{g,m}^n$ if p = m. In particular,

$$\mathbf{1}\bar{*}_{q,m}^{n}u = u \quad \text{for } u \in V. \tag{2.4}$$

Note that if g = 1, $*_{1,m,p}^n$ is the same as $*_{m,p}^n$ defined in [3]. And $*_{g,n,n}^n$ is, in fact, the product $*_{g,n}$ defined in [8]; in particular, $*_{g,0}$ is the product $*_g$ defined in [7].

For $m, n \in (1/T)\mathbb{Z}_+$, let

$$O'_{q,n,m}(V) = \text{span}\{u \circ_{q,m}^n v \mid u, v \in V\} + L_{n,m}(V),$$

where $L_{n,m}(V) = \text{span}\{(L(-1) + L(0) + m - n)u \mid u \in V\}$ and for $u \in V^r, v \in V$,

$$u \circ_{g,m}^{n} v = \operatorname{Res}_{z} \frac{(1+z)^{\operatorname{wt} u - 1 + \delta_{\bar{m}}(r) + \lfloor m \rfloor + r/T}}{z^{\lfloor m \rfloor + \lfloor n \rfloor + \delta_{\bar{m}}(r) + \delta_{\bar{n}}(T-r) + 1}} Y(u,z)v.$$

Again if m = n, then $u \circ_{g,n}^n v = u \circ_{g,n} v$ has been defined in [8]. Then, $O'_{g,n,n}(V) = O_{g,n}(V)$, $O_{g,0}(V) = O_g(V)$, $A_{g,n}(V) = V/O_{g,n}(V)$ and $A_{g,0}(V) = A_g(V)$ (cf. [7,8]).

Lemma 2.4. For any weak g-twisted V-module M, $m, n \in (1/T)\mathbb{Z}_+$ and $a \in O'_{g,n,m}(V)$, we have $o_{m-n}(a) = 0$ on $\Omega_m(M)$.

Proof. It is trivial if a = L(-1)u + (L(0) + m - n)u for some $u \in V$. Assume that a has the form $u \circ_{g,m}^n v$ for some $u \in V^r$ and $v \in V$. Then for any $w \in \Omega_m(M)$, by the twisted Jacobi identity (2.1) we have

$$\begin{split} o_{m-n}\left(u\circ_{g,m}^{n}v\right)w\\ &=\sum_{k\geq0}\left(^{\operatorname{wt}\,u+\lfloor m\rfloor+\delta_{\bar{n}}(T-r)+r/T-1}\right)o_{m-n}\left(u_{k-\lfloor m\rfloor-\lfloor n\rfloor-\delta_{\bar{m}}(r)-\delta_{\bar{n}}(T-r)-1}v\right)w\\ &=\sum_{k\geq0}(-1)^{k}\left(^{-\lfloor m\rfloor-\lfloor n\rfloor-\delta_{\bar{m}}(r)-\delta_{\bar{n}}(T-r)-1}\right)\times\\ &\left(u_{\operatorname{wt}\,u+r/T-\lfloor n\rfloor-\delta_{\bar{n}}(T-r)-2-k}v_{\operatorname{wt}\,v-1+m+k+1+\delta_{\bar{n}}(T-r)-r/T-\bar{n}/T}-\right.\\ &\left.\left(-1\right)^{\lfloor m\rfloor+\lfloor n\rfloor+\delta_{\bar{m}}(r)+\delta_{\bar{n}}(T-r)+1}\right.\\ &\left.\times v_{\operatorname{wt}\,v+\bar{m}/T-r/T-\delta_{\bar{m}}(r)-1-k}u_{\operatorname{wt}\,u-1+m+k+r/T+\delta_{\bar{m}}(r)-\bar{m}/T}\right)w. \end{split}$$

But by the definition of $\Omega_m(M)$,

$$v_{\text{wt}\, v-1+m+k+1+\delta_{\bar{n}}(T-r)-r/T-\bar{n}/T}w = u_{\text{wt}\, u-1+m+k+r/T+\delta_{\bar{m}}(r)-\bar{m}/T}w = 0 \ \text{ for all } k \in \mathbb{Z}_+.$$

Thus, $o_{m-n}\left(u\circ_{q,m}^nv\right)=0$ on $\Omega_m(M)$, completing the proof. \square

The following theorem is from [8, Theorem 2.4 and Theorem 3.3].

Theorem 2.5. (1) The product $\bar{*}_{g,n}^n = *_{g,n} = *_{g,n}^n$ induces the structure of an associative algebra on $A_{g,n}(V)$ with identity $\mathbf{1} + O_{g,n}(V)$.

(2) Suppose that M is a weak g-twisted V-module. Then there is a representation of the associative algebra $A_{g,n}(V)$ on $\Omega_n(M)$ induced by the map $a \mapsto o(a) = a_{\operatorname{wt} a-1}$ for $a \in V$. Moreover, if $M = \bigoplus_{k \in (1/T)\mathbb{Z}_+} M(k)$ is an admissible g-twisted V-module, then $\bigoplus_{0 \leq k \leq n} M(k) \subseteq \Omega_n(M)$ and for each $k \in (1/T)\mathbb{Z}$ such that $0 \leq k \leq n$, M(k) is an $A_{g,n}(V)$ -module.

For any $a, b, c, u \in V$ and any $p_1, p_2, p_3 \in (1/T)\mathbb{Z}_+$, let $O''_{g,n,m}(V)$ be the linear span of

$$u *_{g,m,p_3}^n ((a *_{g,p_1,p_2}^{p_3} b) *_{g,m,p_1}^{p_3} c - a *_{g,m,p_2}^{p_3} (b *_{g,m,p_1}^{p_2} c)).$$

In particular, by (2.4) we have

$$(a *_{g,p_1,p_2}^n b) *_{g,m,p_1}^n c - a *_{g,m,p_2}^n (b *_{g,m,p_1}^{p_2} c) \in O''_{g,n,m}(V).$$
 (2.5)

Let

$$O_{g,n,m}^{\prime\prime\prime}(V) = \sum_{p_1, p_2 \in (1/T)\mathbb{Z}_+} \left(V *_{g,p_1,p_2}^n O_{g,p_2,p_1}^\prime(V) \right) *_{g,m,p_1}^n V$$

and

$$O_{g,n,m}(V) = O'_{g,n,m}(V) + O''_{g,n,m}(V) + O'''_{g,n,m}(V).$$

Set

$$A_{q,n,m}(V) = V/O_{q,n,m}(V).$$

Theorem 2.6. [4] Let V be a vertex operator algebra and $m, n \in (1/T)\mathbb{Z}_+$. Then $A_{g,n,m}(V)$ is an $A_{g,n}(V)-A_{g,m}(V)$ -bimodule such that the left and right actions of $A_{g,n}(V)$ and $A_{g,m}(V)$ are induced by $\bar{*}_{g,m}^n$ and $*_{g,m}^n$, respectively.

Let U be an $A_{q,m}(V)$ -module. Set

$$M(U) = \bigoplus_{n \in (1/T)\mathbb{Z}_+} A_{g,n,m}(V) \otimes_{A_{g,m}(V)} U.$$

Then, M(U) is $(1/T)\mathbb{Z}_+$ -graded such that $M(U)(n) = A_{g,n,m}(V) \otimes_{A_{g,m}(V)} U$. For $u \in V^r$, $p \in r/T + \mathbb{Z}$ and $n \in (1/T)\mathbb{Z}$, define an operator u_p from M(U)(n) to M(U)(n + wt u - p - 1) (with the convention that M(U)(i) = 0 if i < 0) by

$$u_p((v + O_{g,n,m}(V) \otimes w)) = \begin{cases} (u *_{g,m,n}^{\text{wt } u - p - 1 + n} v + O_{g,\text{wt } u - p - 1 + n,m}(V)) \otimes w, & \text{if wt } u - 1 - p + n \ge 0, \\ 0, & \text{if wt } u - 1 - p + n < 0, \end{cases}$$

for $v \in V$ and $w \in U$.

Theorem 2.7. [4] Let U be an $A_{g,m}(V)$ -module. Then

$$M(U) = \bigoplus_{n \in (1/T)\mathbb{Z}_+} A_{g,n,m}(V) \otimes_{A_{g,m}(V)} U$$

is an admissible g-twisted V-module with $M(U)(n) = A_{g,n,m}(V) \otimes_{A_{g,m}(V)} U$ and has the following universal property: for any weak g-twisted V-module W and any $A_{g,m}(V)$ -homomorphism $\phi: U \to \Omega_m(W)$, there is a unique homomorphism $\bar{\phi}: M(U) \to W$ of weak g-twisted V-modules which extends ϕ . Moreover, if U cannot factor through $A_{g,m-1/T}(V)$, then $M(U)(0) \neq 0$.

3. Associative algebras $\mathcal{A}_{g,n}(V)$ and $\mathcal{A}_{g,n}(V)-\mathcal{A}_{g,m}(V)$ -bimodules $\mathcal{A}_{g,n,m}(V)$

In this section, we will construct a family of associative algebras $\mathcal{A}_{g,n}(V)$ and a family of $\mathcal{A}_{g,n}(V) - \mathcal{A}_{g,m}(V)$ -bimodules $\mathcal{A}_{g,n,m}(V)$ from the perspective of representations, and show that they share the similar properties as for $A_{g,m}(V)$ and $A_{g,n,m}(V)$.

For any $m, n \in (1/T)\mathbb{Z}_+$, let

$$\mathcal{O}_{g,n,m}(V) = \left\{u \in V \mid o_{m-n}(u)|_{\Omega_m(M)} = 0 \text{ for any weak } g\text{-twisted } V\text{-module } M\right\}.$$

Set

$$\mathcal{A}_{g,n,m}(V) = V/\mathcal{O}_{g,n,m}(V).$$

Write $\mathcal{O}_{g,n,m}(V)$ as $\mathcal{O}_{g,n}(V)$ and $\mathcal{A}_{g,n,m}(V)$ as $\mathcal{A}_{g,n}(V)$ if m=n. And when g=1, these $\mathcal{A}_{1,n,m}(V)$ were studied in [15].

The following lemma is clear from the definition of $\mathcal{O}_{q,n,m}(V)$.

Lemma 3.1. $\mathcal{O}_{g,n,m}(V) \subseteq \mathcal{O}_{g,n-l,m-l}(V)$ for $(1/T)\mathbb{Z}_+ \ni l \leq \min\{m,n\}$.

We have the following result from [4, Lemma 5.1].

Lemma 3.2. Let M be a weak q-twisted V-module. Then

$$o_{m-n}\left(u*_{q,m,p}^nv\right)=o_{p-n}(u)o_{m-p}(v) \ on \ \Omega_m(M) \ for \ u,v\in V \ and \ m,n,p\in (1/T)\mathbb{Z}_+.$$

In particular, $u *_{q,m}^{n} \mathbf{1} - u \in \mathcal{O}_{g,n,m}(V)$.

By the definition of $\mathcal{O}_{g,n,m}(V)$, Lemma 2.3 and Lemma 3.2, it is not difficult to show the following statements.

Lemma 3.3. Let $m, n \in (1/T)\mathbb{Z}_+$. Then

- $(1) \left(a *_{g,p_1,p_2}^n b\right) *_{g,m,p_1}^n c a *_{g,m,p_2}^n \left(b *_{g,m,p_1}^{p_2} c\right) \in \mathcal{O}_{g,n,m}(V) \ \text{for } a,b,c \in V \ \text{and} \\ p_1,p_2 \in (1/T)\mathbb{Z}_+. \ \text{In particular, } \left(a \bar{*}_{g,m}^n b\right) *_{g,m}^n c \equiv a \bar{*}_{g,m}^n \left(b *_{g,m}^n c\right) \ \text{mod} \ \mathcal{O}_{g,n,m}(V).$
- $(2) V *_{g,m,p}^{n} \mathcal{O}_{g,p,m}(V) \subseteq \mathcal{O}_{g,n,m}(V) \text{ and } \mathcal{O}_{g,n,p}(V) *_{g,m,p}^{n} V \subseteq \mathcal{O}_{g,n,m}(V) \text{ for } p \in (1/T)\mathbb{Z}_{+}. \text{ In particular, } V \bar{*}_{g,m}^{n} \mathcal{O}_{g,n,m}(V) \subseteq \mathcal{O}_{g,n,m}(V) \text{ and } \mathcal{O}_{g,n,m}(V) *_{g,m}^{n} V \subseteq \mathcal{O}_{g,n,m}(V); \mathcal{O}_{g,n}(V) \bar{*}_{g,m}^{n} V \subseteq \mathcal{O}_{g,n,m}(V) \text{ and } V *_{g,m}^{n} \mathcal{O}_{g,m}(V) \subseteq \mathcal{O}_{g,n,m}(V).$

Remark 3.4. (1) It is clear by Lemma 2.4, Lemma 3.2 and Lemma 3.3 that $O_{g,n,m}(V) = O'_{g,n,m}(V) + O''_{g,n,m}(V) + O'''_{g,n,m}(V) \subseteq \mathcal{O}_{g,n,m}(V)$. In particular, $O_{g,n}(V) \subseteq \mathcal{O}_{g,n}(V)$.

(2) Since $A_{g,n}(V)$ is a quotient algebra of $A_{g,n}(V)$, every $A_{g,n}(V)$ -module automatically becomes an $A_{g,n}(V)$ -module.

Theorem 3.5. Let $m, n \in (1/T)\mathbb{Z}_+$. Then

- (1) $\mathcal{A}_{g,n}(V)$ is an associative algebra under the multiplication $\bar{*}_{g,n}^n = *_{g,n}^n$ with the identity $\mathbf{1} + \mathcal{O}_{g,n}(V)$.
- (2) $\mathcal{A}_{g,n,m}(V)$ is an $\mathcal{A}_{g,n}(V)-\mathcal{A}_{g,m}(V)$ -bimodule with $\bar{*}_{g,m}^n$ the left action and $*_{g,m}^n$ the right action.
- (3) Suppose that M is a weak g-twisted V-module. Then there is a representation of $\mathcal{A}_{g,n}(V)$ on $\Omega_n(M)$ induced by the linear map $u \mapsto o(u)$ for $u \in V$. Moreover, if $M = \bigoplus_{k \in (1/T)\mathbb{Z}_+} M(k)$ is an admissible g-twisted V-module, then

$$\bigoplus_{0 < k < n} M(k) \subseteq \Omega_n(M)$$

and each M(k) is an $\mathcal{A}_{q,n}(V)$ -module for $0 \leq k \leq n$.

Proof. (1) By Lemma 3.3 (2), $\mathcal{O}_{g,n}(V)$ is a two-sided ideal of V under the multiplication $\bar{*}_{g,n}^n = *_{g,n}^n$. It follows from Lemma 3.3 (1) that this multiplication satisfies the associativity. Thus, $\mathcal{A}_{g,n}(V)$ is an associative algebra. And by (2.4) and Lemma 3.2 we see that $\mathbf{1} + \mathcal{O}_{g,n}(V)$ is its identity.

- (2) Note that the left action of $\mathcal{A}_{g,n}(V)$ and the right action of $\mathcal{A}_{g,m}(V)$ on $\mathcal{A}_{g,n,m}(V)$ are well defined by Lemma 3.3(2); and also that the two actions are compatible by Lemma 3.3(1). Thus, $\mathcal{A}_{g,n,m}(V)$ is an $\mathcal{A}_{g,n}(V)-\mathcal{A}_{g,m}(V)$ -bimodule.
- (3) It is clear from the definition of $\mathcal{O}_{g,n}(V)$ that the given representation is well defined. And this map is an algebra homomorphism by Lemma 3.2, proving the first statement. The second statement follows from that $v_{\text{wt }v-1+i}M(k)=0$ for any i>n and $k\leq n$ and that $o(v)M(k)\subseteq M(k)$ for any $k\in (1/T)\mathbb{Z}_+$. \square

The following corollary is an immediate consequence of Lemma 3.1 and Theorem 3.5(2).

Corollary 3.6. For any $l, m, n \in (1/T)\mathbb{Z}_+$ such that $l \leq \min\{m, n\}$, the identity map on V induces an epimorphism of $\mathcal{A}_{g,n}(V) - \mathcal{A}_{g,m}(V)$ -bimodules from $\mathcal{A}_{g,n,m}(V)$ to $\mathcal{A}_{g,n-l,m-l}(V)$. In particular, the identity map on V induces an epimorphism of algebras from $\mathcal{A}_{g,n}(V)$ to $\mathcal{A}_{g,n-l}(V)$.

Let U be an $\mathcal{A}_{g,m}(V)$ -module. Set

$$\mathcal{M}(U) = \bigoplus_{n \in (1/T)\mathbb{Z}_{\perp}} \mathcal{A}_{g,n,m}(V) \otimes_{\mathcal{A}_{g,m}(V)} U.$$

Then $\mathcal{M}(U) = \bigoplus_{n \in (1/T)\mathbb{Z}_+} \mathcal{M}(U)(n)$ is $(1/T)\mathbb{Z}_+$ -graded with $\mathcal{M}(U)(n) = \mathcal{A}_{g,n,m}(V)$ $\otimes_{\mathcal{A}_{g,m}(V)} U$ for $n \in (1/T)\mathbb{Z}_+$. Following the construction of M(U) (see Section 2), for $u \in V^r$ define the vertex operator $Y_{\mathcal{M}(U)}(u,z) = \sum_{p \in r/T+\mathbb{Z}} u_p z^{-p-1}$ with u_p being a linear map from $\mathcal{M}(U)(n)$ to $\mathcal{M}(U)(n+\operatorname{wt} u-p-1)$ for $n \in (1/T)\mathbb{Z}_+$ (decreeing $\mathcal{M}(U)(k) = 0$ if k < 0) given by

$$u_p((v + \mathcal{O}_{g,n,m}(V)) \otimes w) = (u *_{g,m,n}^{n+\text{wt } u-p-1} v + \mathcal{O}_{g,m,n+\text{wt } u-p-1}(V)) \otimes w$$

for $v \in V$ and $w \in U$. This action is well defined, since for any $v \in \mathcal{A}_{g,n,m}(V), a \in \mathcal{A}_{g,m}(V)$ and $w \in U$, we have $u *_{g,m,n}^{n+\text{wt}\,u-p-1} \mathcal{O}_{g,n,m}(V) \subseteq \mathcal{O}_{g,n+\text{wt}\,u-p-1,m}(V)$ by Lemma 3.3 (2), and

$$u_p\left(\left(v*_{g,m}^n a\right) \otimes w\right) = \left(u*_{g,m,n}^{n+\operatorname{wt} u-p-1} \left(v*_{g,m}^n a\right)\right) \otimes w$$

$$= \left(\left(u*_{g,m,n}^{n+\operatorname{wt} u-p-1} v\right) *_{g,m}^{n+\operatorname{wt} u-p-1} a\right) \otimes w$$

$$= \left(u*_{g,m,n}^{n+\operatorname{wt} u-p-1} v\right) \otimes a \cdot w = u_p(v \otimes a \cdot w)$$

by Lemma 3.3 (1). Note that U can also be viewed as an $A_{g,m}(V)$ -module by Remark 3.4 (2). Then we can define the linear map

$$\psi_{n,m}: A_{g,n,m}(V) \otimes_{A_{g,m}(V)} U \longrightarrow \mathcal{A}_{g,n,m}(V) \otimes_{\mathcal{A}_{g,m}(V)} U$$

sending $(u + O_{g,n,m}(V)) \otimes w$ to $(u + O_{g,n,m}(V)) \otimes w$ for $u \in V$ and $w \in U$, which is well defined. Note that these $\psi_{n,m}$ for $n \in (1/T)\mathbb{Z}_+$ induce the surjective linear map $\psi: M(U) \longrightarrow \mathcal{M}(U)$ such that $\psi(u_p w) = u_p \psi(w)$ for any $p \in (1/T)\mathbb{Z}, u \in V$ and $w \in M(U)$. Thus, $\mathcal{M}(U)$ is a weak g-twisted V-module. Moreover, by definition of the action of u_p , $\mathcal{M}(U)$ is an admissible g-twisted V-module. Similarly, one can show that $\mathcal{M}(U)$ shares the same universal property as M(U). Then we arrive at the following result.

Theorem 3.7. Let U be an $A_{g,m}(V)$ -module. Then

$$\mathcal{M}(U) = \bigoplus_{n \in (1/T)\mathbb{Z}_+} \mathcal{A}_{g,n,m}(V) \otimes_{\mathcal{A}_{g,m}(V)} U$$

is an admissible g-twisted V-module with $\mathcal{M}(U)(n) = \mathcal{A}_{g,n,m}(V) \otimes_{\mathcal{A}_{g,m}(V)} U$ satisfying the following universal property: for any weak g-twisted V-module W and any $\mathcal{A}_{g,m}(V)$ -morphism $\phi: U \to \Omega_m(W)$, there is a unique homomorphism $\bar{\phi}: \mathcal{M}(U) \to W$ of weak g-twisted V-modules which extends ϕ . Moreover, $\mathcal{M}(U)(0) \neq 0$ if U cannot factor through $\mathcal{A}_{g,m-1/T}(V)$.

4. The universal enveloping algebra U(V[g]) of V with respect to g

In this section, we shall first recall the construction of the universal enveloping algebra U(V[g]) of V with respect to g and then use U(V[g]) to construct admissible g-twisted V-modules with some universal property.

Recall from [7] (see also [1]) the Lie algebra

$$\hat{V}[g] = \mathcal{L}(V, g) / D\mathcal{L}(V, g),$$

where $\mathcal{L}(V,g) = \bigoplus_{r=0}^{T-1} V^r \otimes \mathbb{C}t^{\frac{r}{T}} \left[t,t^{-1}\right]$ and $D = L(-1) \otimes \mathrm{id} + \mathrm{id} \otimes \frac{d}{dt}$. Denote by u(m) the image of $u \otimes t^m$ in $\hat{V}[g]$. Then the Lie bracket on $\hat{V}[g]$ is given by

$$\left[u(m+\frac{r}{T}),v(n+\frac{s}{T})\right] = \sum_{i=0}^{\infty} {m+\frac{r}{T} \choose i} (u_i v)(m+n+\frac{r+s}{T}-i)$$

for $u \in V^r, v \in V^s$ and $m, n \in \mathbb{Z}$. If we define the degree of u(m) to be wt u-m-1, then $\hat{V}[g]$ is a $(1/T)\mathbb{Z}$ -graded Lie algebra, i.e., $\hat{V}[g] = \bigoplus_{m \in (1/T)\mathbb{Z}} \hat{V}[g]_m$ and $\left[\hat{V}[g]_i, \hat{V}[g]_j\right] \subseteq \hat{V}[g]_{i+j}$ for any $i, j \in (1/T)\mathbb{Z}$.

Let $U(\hat{V}[g])$ be the universal enveloping algebra of the Lie algebra $\hat{V}[g]$. Then the $(1/T)\mathbb{Z}$ -grading on $\hat{V}[g]$ induces a $(1/T)\mathbb{Z}$ -grading on $U(\hat{V}[g]) = \bigoplus_{m \in (1/T)\mathbb{Z}} U(\hat{V}[g])_m$. Set

$$U(\hat{V}[g])_{m}^{k} = \sum_{i \leq k, i \in (1/T)\mathbb{Z}} U(\hat{V}[g])_{m-i} U(\hat{V}[g])_{i}$$

for $0 > k \in (1/T)\mathbb{Z}$ and $U(\hat{V}[g])_m^0 = U(\hat{V}[g])_m$. Then

$$U(\hat{V}[g])_m^k \subseteq U(\hat{V}[g])_m^{k+1/T}$$

and

$$\bigcap_{k \in -(1/T)\mathbb{Z}_+} U(\hat{V}[g])_m^k = 0, \quad \bigcup_{k \in -(1/T)\mathbb{Z}_+} U(\hat{V}[g])_m^k = U(\hat{V}[g])_m.$$

Thus, $\left\{U(\hat{V}[g])_m^k\mid k\in -(1/T)\mathbb{Z}_+\right\}$ forms a fundamental neighborhood system of $U(\hat{V}[g])_m$. Let $\tilde{U}(\hat{V}[g])_m$ be the completion of $U(\hat{V}[g])_m$, then

$$\widetilde{U}(\widehat{V}[g]) := \bigoplus_{m \in (1/T)\mathbb{Z}} \widetilde{U}(\widehat{V}[g])_m$$

is a complete topological ring which allows infinite sums in it.

For each $m \in (1/T)\mathbb{Z}$, define a linear map $J_m(\cdot): V \to \hat{V}[g]$ sending $u \in V^r$ to $u(\operatorname{wt} u + m - 1)$ if $m \in r/T + \mathbb{Z}$ and zero otherwise.

Definition 4.1. The universal enveloping algebra U(V[g]) of V with respect to g is the quotient of $\widetilde{U}(\hat{V}[g])$ by a two-sided ideal generated by the following relations:

$$\mathbf{1}(i) = \delta_{i,-1} \text{ for } i \in \mathbb{Z},$$
$$[\omega(i+1), \omega(j+1)] = (i-j)\omega(i+j+1) + \delta_{i+j,0} \frac{i^3 - i}{12}c \quad \text{ for } i, j \in \mathbb{Z},$$

and

$$\sum_{i\geq 0} (-1)^i \binom{l}{i} \left(J_{s-i}(u) J_{t+i}(v) - (-1)^l J_{l+t-i}(v) J_{s+i-l}(u) \right)$$

$$= \sum_{i\geq 0} \binom{s + \operatorname{wt} u - l - 1}{i} J_{s+t}(u_{l+i}v) \quad \text{for } u \in V^r, v \in V^{r'}, l \in \mathbb{Z}, s \in \frac{r}{T} + \mathbb{Z}, t \in \frac{r'}{T} + \mathbb{Z}.$$

It is clear that $U(V[g])=\bigoplus_{m\in(1/T)\mathbb{Z}}U(V[g])_m$ is a $(1/T)\mathbb{Z}$ -graded associative algebra. Set

$$U(V[g])_{m}^{k} = \sum_{i < k, i \in (1/T)\mathbb{Z}} U(V[g])_{m-i} U(V[g])_{i}$$

for $0 > k \in (1/T)\mathbb{Z}$. Then, $U(V[g])_{n-m}/U(V[g])_{n-m}^{-m-1/T}$ is a $U(V[g])_0/U(V[g])_0^{-n-1/T} - U(V[g])_0/U(V[g])_0^{-m-1/T}$ -bimodule.

Theorem 4.2. [16] The linear map $\varphi: V \to U(V[g])_0$ sending u to $J_0(u)$ induces an algebra isomorphism φ_n between $A_{g,n}(V)$ and $U(V[g])_0/U(V[g])_0^{-n-1/T}$ for each $n \in (1/T)\mathbb{Z}_+$.

Remark 4.3. (1) Suppose that M is a weak g-twisted V-module. Then by Theorem 4.2 and Theorem 2.5 (2), there is a representation of $U(V[g])_0/U(V[g])_0^{-n-1/T}$ on $\Omega_n(M)$ induced by the linear map $a \to o(a) = a_{\text{wt } a-1}$ for $a \in V$.

(2) From the construction of U(V[g]), any weak g-twisted V-module is naturally a U(V[g])-module with the action induced by the map $u(m) \mapsto u_m$ for any $u \in V^r$ and $m \in r/T + \mathbb{Z}$.

Theorem 4.2 tells us that $A_{g,n}(V)$ can be realized as some quotient of $U(V[g])_0$. And in Section 5 we are going to make use of the subspace $U(V[g])_{n-m}$ to realize the bimodule $A_{g,n,m}(V)$.

Let U be a $U(V[g])_0/U(V[g])_0^{-m-1/T}$ -module. Set

$$\mathbb{M}(U) = \bigoplus_{n \in (1/T)\mathbb{Z}_+} U(V[g])_{n-m}/U(V[g])_{n-m}^{-m-1/T} \otimes_{U(V[g])_0/U(V[g])_0^{-m-1/T}} U.$$

Set $\mathbb{M}(U) = \bigoplus_{n \in (1/T)\mathbb{Z}_+} \mathbb{M}(U)(n)$ with

$$\mathbb{M}(U)(n) = U(V[g])_{n-m}/U(V[g])_{n-m}^{-m-1/T} \otimes_{U(V[g])_0/U(V[g])_0^{-m-1/T}} U$$

for $n \in (1/T)\mathbb{Z}_+$. Then, $\mathbb{M}(U)(m) \cong U$ as $A_{g,m}(V)$ -modules by Theorem 4.2.

Also following the construction of M(U), we equip $\mathbb{M}(U)$ with the vertex operator maps $Y_{\mathbb{M}(U)}(u,z) = \sum_{p \in r/T+\mathbb{Z}} u_p z^{-p-1}$ for $u \in V^r$, where for $n \in (1/T)\mathbb{Z}_+$, the linear map u_p from $\mathbb{M}(U)(n)$ to $\mathbb{M}(U)(n+\operatorname{wt} u-p-1)$ is defined as follows:

$$u_p(v \otimes w) = \begin{cases} u(p)v \otimes w, & \text{if } n + \text{wt } u - p - 1 \ge 0, \\ 0, & \text{if } n + \text{wt } u - p - 1 < 0, \end{cases}$$

for $v \in U(V[g])_{n-m}/U(V[g])_{n-m}^{-m-1/T}$ and $w \in U$. Then $\mathbb{M}(U)$ is an admissible g-twisted V-module, since the twisted Jacobi identity follows immediately from the construction of U(V[g]); and $\mathbb{M}(U)$ is generated by $\mathbb{M}(U)(m)$.

Theorem 4.4. Let U be a $U(V[g])_0/U(V[g])_0^{-m-1/T}$ -module. Then the admissible g-twisted V-module

$$\mathbb{M}(U) = \bigoplus_{n \in (1/T)\mathbb{Z}_+} U(V[g])_{n-m} / U(V[g])_{n-m}^{-m-1/T} \otimes_{U(V[g])_0 / U(V[g])_0^{-m-1/T}} U$$

has the following universal property: for any weak g-twisted V-module W and any $A_{g,m}(V)$ -morphism $\phi: U \to \Omega_m(W)$, there is a unique homomorphism $\bar{\phi}: \mathbb{M}(U) \to W$ of weak g-twisted V-modules which extends ϕ . Moreover, if U cannot factor through $A_{g,m-1/T}(V)$, then $\mathbb{M}(U)(0) \neq 0$.

Proof. Define $\bar{\phi}: \mathbb{M}(U) \to W$ by $\bar{\phi}(u \otimes w) = u\phi(w)$ for $u \in U(V[g])_{n-m}/U(V[g])_{n-m}^{-m-1/T}$ and $w \in U$. Note that the action of $U(V[g])_{n-m}^{-m-1/T}$ on $\Omega_m(W)$ is trivial by Remark 4.3 (2) and also that

$$\bar{\phi}(u\cdot v\otimes w)=(u\cdot v)\phi(w)=u\phi(v\cdot w)=\bar{\phi}(u\otimes v\cdot w)\quad\text{for }v\in U(V[g])_0/U(V[g])_0^{-m-1/T}.$$

Thus, $\bar{\phi}$ is well defined. It is clear that $\bar{\phi}|_{U} = \phi$ by regarding

$$U = \mathbb{M}(U)(m) = U(V[g])_0 / U(V[g])_0^{-m-1/T} \otimes_{U(V[g])_0 / U(V[g])_0^{-m-1/T}} U.$$

And for $v \in V$, again by Remark 4.3(2), we have

$$\bar{\phi}(v_p(u \otimes w)) = \bar{\phi}(v(p)u \otimes v) = (v(p)u)\phi(w) = v(p)(u\phi(w)) = v(p)(\bar{\phi}(u \otimes w))$$
$$= v_p(\bar{\phi}(u \otimes w)).$$

Thus, $\bar{\phi}$ is a homomorphism of weak g-twisted V-modules, whose uniqueness follows from the fact that $\mathbb{M}(U)$ is generated by $U = \mathbb{M}(U)(m)$. \square

5. Isomorphisms

In this section we shall show $O_{g,n,m}(V) = \mathcal{O}_{g,n,m}(V)$ and realize $A_{g,n,m}(V)$ as some quotient of $U(V[g])_{n-m}$.

Following from [4, Lemma 5.1], we have:

Lemma 5.1. Let $u, v \in V$ and $m, n, p \in (1/T)\mathbb{Z}_+$. Then

$$J_{m-n}\left(u *_{q,m,p}^{n} v\right) \equiv J_{p-n}(u)J_{m-p}(v) \mod U(V[g])_{n-m}^{-m-1/T}.$$

The main result of this section is as follows.

Theorem 5.2. (1) $A_{g,m}(V) = A_{g,m}(V) = A_{g,m,m}(V)$ and $A_{g,n,m}(V) = A_{g,n,m}(V)$ for any $m, n \in (1/T)\mathbb{Z}_+$.

(2) The $A_{g,n}(V)-A_{g,m}(V)$ -bimodules $A_{g,n,m}(V)$ and $U(V[g])_{n-m}/U(V[g])_{n-m}^{-m-1/T}$ are isomorphic for any $m, n \in (1/T)\mathbb{Z}_+$.

Proof. Fix $m \in (1/T)\mathbb{Z}_+$ and take $U = A_{q,m}(V)$. By Theorem 4.2,

$$A_{g,m}(V) \cong \mathbb{M}(U)(m) = U(V[g])_0/U(V[g])_0^{-m-1/T} \otimes_{U(V[g])_0/U(V[g])_0^{-m-1/T}} A_{g,m}(V).$$

Now it follows from Theorem 3.5 (3) that the multiplication $\bar{*}_{g,m}^m = *_{g,m}^m$ on V induces an $\mathcal{A}_{g,m}(V)$ -module structure on $A_{g,m}(V)$. In particular, we have $\mathcal{O}_{g,m}(V) \subseteq O_{g,m}(V)$, which together with Remark 3.4 (1) gives $\mathcal{O}_{g,m}(V) = O_{g,m}(V)$, i.e., $\mathcal{A}_{g,m}(V) = A_{g,m}(V)$. Similarly, when replacing $\mathbb{M}(U)$ by M(U) we can obtain $A_{g,m,m}(V) = \mathcal{A}_{g,m}(V)$. That is, both M(U) and $\mathcal{M}(U)$ have the same generating set. Then $M(U) = \mathcal{M}(U)$, since these two modules have the same universal property by Theorems 2.7 and 3.7. Now by considering M(U)(n) and $\mathcal{M}(U)(n)$ we see that $A_{g,n,m}(V) = \mathcal{A}_{g,n,m}(V)$, proving (1).

For convenience, we would identify $A_{g,n,m}(V) \otimes_{A_{g,m}(V)} A_{g,m}(V)$ with $A_{g,n,m}(V)$, and identify $U(V[g])_{n-m}^{-m-1/T} \otimes_{U(V[g])_0/U(V[g])_0^{-m-1/T}} A_{g,m}(V)$ with $U(V[g])_{n-m}^{-m-1/T}$. It follows from (1) and Theorem 4.2 that the linear map

$$\varphi_{m,m}: A_{g,m,m}(V) \to U(V[g])_0/U(V[g])_0^{-m-1/T}$$

 $u + O_{g,m,m}(V) \mapsto J_0(u) + U(V[g])_0^{-m-1/T}$

is an isomorphism of $A_{g,m}(V)$ -modules. Now by Theorem 2.7 and Theorem 4.4, this map can be extended to an isomorphism of admissible g-twisted V-modules from M(U) to $\mathbb{M}(U)$ such that for any $n \in (1/T)\mathbb{Z}_+$, the linear map

$$\varphi_{n,m}: A_{g,n,m}(V) \to U(V[g])_{n-m}/U(V[g])_{n-m}^{-m-1/T}$$

 $u + O_{g,n,m}(V) \mapsto J_{m-n}(u) + U(V[g])_{n-m}^{-m-1/T}$

gives an isomorphism of $A_{g,n}(V)$ -modules. In fact, this is also a homomorphism of right $A_{g,m}(V)$ -modules:

$$\begin{split} & \varphi_{n,m} \left((u + O_{g,n,m}(V)) *_{g,m}^m \left(b + O_{g,m}(V) \right) \right) = \varphi_{n,m} \left(u *_{g,m}^n b + O_{g,n,m}(V) \right) \\ = & J_{m-n} \left(u *_{g,m}^n b \right) + U(V[g])_{n-m}^{-m-1/T} = J_{m-n}(u) J_0(b) + U(V[g])_{n-m}^{-m-1/T} \\ = & \left(J_{m-n}(u) + U(V[g])_{n-m}^{-m-1/T} \right) \cdot (b + O_{g,m}(V)) \\ = & \varphi_{n,m} \left(u + O_{g,n,m}(V) \right) \cdot (b + O_{g,m}(V)) \,, \end{split}$$

where the third equality follows from Lemma 5.1. Thus $\varphi_{n,m}$ is an $A_{g,n}(V)-A_{g,m}(V)$ -bimodule isomorphism, proving (2). \square

By Theorem 5.2 and Corollary 3.6, one can obtain the following result.

Corollary 5.3. (1) The identity map on V induces an epimorphism of $A_{g,n}(V) - A_{g,m}(V)$ -bimodules from $A_{g,n,m}(V)$ to $A_{g,n-l,m-l}(V)$ for $l \in (1/T)\mathbb{Z}_+$ such that $l \leq \min\{m,n\}$. (2) $O_{g,n,n}(V) = O_{g,n}(V)$ and

$$O_{g,n,m}(V) = \left\{ u \in V \mid J_{m-n}(u) \in U(V[g])_{n-m}^{-m-1/T} \right\}$$
$$= \left\{ u \in V \mid o_{m-n}(u) \mid_{\Omega_m(M)} = 0 \text{ for any weak } g\text{-twisted } V\text{-module } M \right\}$$

Remark 5.4. The equality $A_{g,n,n}(V) = A_{g,n}(V)$ and Corollary 5.3 (1) were first proved in [4].

6. Refining bimodules

In this section we shall refine the definition of the $A_{g,n}(V)-A_{g,m}(V)$ -bimodule $A_{g,n,m}(V)$.

Notation 6.1. Until further notice we shall use the following conventions.

- (1) For $m \in (1/T)\mathbb{Z}$ and $i \in \mathbb{Z}$, $\binom{m}{i} = 1$ if i = 0, and 0 if i < 0.
- (2) For $k, l \in (1/T)\mathbb{Z}$, define $\sum_{i=k}^{l} a_i = \sum_{i \in \mathbb{Z}_{k,l}} a_i$, where $\mathbb{Z}_{k,l} = \begin{cases} \mathbb{Z} \cap [l,k] & \text{if } l \leq k, \\ \mathbb{Z} \cap [k,l] & \text{if } l > k. \end{cases}$
- (3) For $n \in (1/T)\mathbb{Z}_+$, $a \in V^r$ and $b \in V$, denote $f_i(a,b)$ as follows:

$$f_i(a,b) = \frac{(1+z)^{\text{wt } a+q}}{z^i} Y(a,z)b \text{ for } i \in \mathbb{Z},$$

where $q = -1 + \lfloor n \rfloor + \delta_{\bar{n}}(r) + r/T$.

Lemma 6.2. Let $n \in (1/T)\mathbb{Z}$ and $l \in \mathbb{Z}$. Then

$$\sum_{i=0}^{n+1+l} (-1)^j \binom{l}{j} \sum_{i=0}^{n+1+l-j} (-1)^i \binom{-l+i+j-1}{i} \frac{1}{z^{i+j}} = 1.$$

Proof. This formula follows from [15, Lemma 3.8] if $n+1+l \geq 0$ and Notation 6.1 (2) if n+1+l < 0. \square

Lemma 6.3. For $k, n \in (1/T)\mathbb{Z}_+$, $a \in V^r, b \in V$ and $j, l \in \mathbb{Z}$, we have

$$a *_{g,n,k+1+q+l-j}^{k} b = \sum_{i=0}^{k+1+q+l-j} (-1)^{i} {\binom{-l+i+j-1}{i}} \operatorname{Res}_{z} f_{i+j-l}(a,b),$$

where $q = -1 + \lfloor n \rfloor + \delta_{\bar{n}}(r) + r/T$.

Proof. Note that $\lfloor k+1+q+l-j\rfloor = \lfloor n\rfloor + \lfloor k+r/T\rfloor + \delta_{\bar{n}}(r) + l-j$ and $\lfloor k\rfloor - \lfloor k+r/T\rfloor + \delta_{\bar{k}}(T-r) = 0$. Then this lemma follows from the definition of product $*_{g,m,p}^n$ and Notation 6.1 (2)-(3). \square

Set

$$M_g^{(m)} = \bigoplus_{n \in (1/T)\mathbb{Z}_+} V/O_{g,n,m}''(V),$$

which is clearly $(1/T)\mathbb{Z}_+$ -graded with $M_g^{(m)}(n) = V/O_{g,n,m}''(V)$. For $u \in V$ and $p \in (1/T)\mathbb{Z}$, define the vertex operator map

$$u_p(v + O_{g,n,m}''(V)) = \begin{cases} u *_{g,m,n}^{n+\operatorname{wt} u - p - 1} v + O_{g,n+\operatorname{wt} u - p - 1,m}'(V), & \text{if } n + \operatorname{wt} u - p - 1 \ge 0, \\ 0, & \text{otherwise.} \end{cases}$$

This action is well defined by the proof in [4, Lemma 3.8].

Lemma 6.4. Let $m, n \in (1/T)\mathbb{Z}_+$. Then

- (1) for any $u \in V^r$ and $p \in r/T + \mathbb{Z}$, $u_p(M_g^{(m)}(n)) = 0$ if p > wt u + n 1;
- (2) $Y_{M_a^{(m)}}(\mathbf{1}, z) = id;$
- (3) for any $a \in V^r$ and $b \in V^s$, we have

$$\left(z_{2}+z_{0}\right)^{\text{wt }a+q}Y_{M_{q}^{(m)}}\left(Y\left(a,z_{0}\right)b,z_{2}\right)\!=\!\left(z_{0}+z_{2}\right)^{\text{wt }a+q}Y_{M_{q}^{(m)}}\left(a,z_{0}+z_{2}\right)Y_{M_{q}^{(m)}}\left(b,z_{2}\right)$$

or equivalently, for any $l \in \mathbb{Z}$,

$$\begin{split} &\operatorname{Res}_{z_{0}} z_{0}^{l} \left(z_{2} + z_{0}\right)^{\operatorname{wt} a + q} z_{2}^{\operatorname{wt} b - q} Y_{M_{g}^{(m)}} \left(Y\left(a, z_{0}\right) b, z_{2}\right) \\ &= \operatorname{Res}_{z_{0}} z_{0}^{l} \left(z_{0} + z_{2}\right)^{\operatorname{wt} a + q} z_{2}^{\operatorname{wt} b - q} Y_{M_{g}^{(m)}} \left(a, z_{0} + z_{2}\right) Y_{M_{g}^{(m)}} \left(b, z_{2}\right) \end{split}$$

on $M_g^{(m)}(n)$, where $q = -1 + \lfloor n \rfloor + \delta_{\bar{n}}(r) + r/T$.

Proof. (1) follows immediately from the definition of u_p . And for (2), it is sufficient to show $\mathbf{1}_p = \delta_{p,-1}$ id on $M_g^{(m)}(n)$ for any $n \in (1/T)\mathbb{Z}_+$. By (1), $\mathbf{1}_p = 0$ on $M_g^{(m)}(n)$ if p > n - 1. Now considering $\mathbb{Z} \ni p \le n - 1$, then for any $v \in V$, we have

$$\mathbf{1}_{p}(v + O_{g,n,m}''(V)) = \mathbf{1} *_{g,m,n}^{n-p-1} v + O_{g,n-p-1,m}''(V)$$

$$= \sum_{i=0}^{\lfloor n \rfloor} (-1)^{i} \binom{\lfloor m \rfloor + \lfloor n-p-1 \rfloor - \lfloor n \rfloor + i}{i}$$

$$\cdot \operatorname{Res}_{z} \frac{(1+z)^{\lfloor m \rfloor}}{z^{\lfloor m \rfloor + \lfloor n-p-1 \rfloor - \lfloor n \rfloor + i + 1}} Y(\mathbf{1}, z) v + O_{g,n-p-1,m}''(V)$$

$$\begin{split} &=\sum_{i=0}^{\lfloor n\rfloor}(-1)^i\binom{\lfloor m\rfloor+\lfloor n-p-1\rfloor-\lfloor n\rfloor+i}{i}\\ &\cdot\binom{\lfloor m\rfloor}{\lfloor m\rfloor+\lfloor n-p-1\rfloor-\lfloor n\rfloor+i}v+O''_{g,n-p-1,m}(V)\\ &=\sum_{i=0}^{\lfloor n\rfloor}(-1)^i\binom{\lfloor m\rfloor-p+i-1}{i}\binom{\lfloor m\rfloor}{\lfloor m\rfloor-p+i-1}v+O''_{g,n-p-1,m}(V)\\ &=\sum_{i=0}^{\lfloor n\rfloor}(-1)^i\binom{\lfloor m\rfloor}{p+1}\binom{p+1}{i}v+O''_{g,n-p-1,m}(V)\\ &=\delta_{p,-1}v+O''_{g,n-p-1,m}(V) & \text{(by Notation 6.1 (1))}\\ &=\delta_{p,-1}(v+O''_{g,n,m}(V)). \end{split}$$

Thus, (2) holds.

The idea of the proof of the third statement comes essentially from [4, Lemma 5.10] (see also [15, Lemma 3.10]). For $v + O''_{g,n,m}(V) \in M_g^{(m)}(n)$, $q = -1 + \lfloor n \rfloor + \delta_{\bar{n}}(r) + r/T$ and let $\alpha \in \{0, \ldots, T-1\}$ be such that $\alpha \equiv \bar{n} - r - s \mod T$, we have

$$\begin{split} &\operatorname{Res}_{z_0} z_0^l \left(z_2 + z_0\right)^{\operatorname{wt} a + q} z_2^{\operatorname{wt} b - q} Y_{M_g^{(m)}} \left(Y\left(a, z_0\right) b, z_2\right) \left(v + O_{g,n,m}''(V)\right) \\ &= \sum_{j \in \mathbb{Z}_+} \left(\overset{\text{wt} a + q}{j} \right) z_2^{\operatorname{wt} a + \operatorname{wt} b - j} Y_{M_g^{(m)}} \left(a_{j+l} b, z_2\right) \left(v + O_{g,n,m}''(V)\right) \\ &= \sum_{j \in \mathbb{Z}_+} \left(\overset{\text{wt} a + q}{j} \right) \sum_{k \in \frac{m}{2} + \mathbb{Z}_+} z_2^{l+k-n+1} \left(a_{j+l} b\right)_{\operatorname{wt} a + \operatorname{wt} b - j - l - 2 - k + n} \left(v + O_{g,n,m}''(V)\right) \\ &= \sum_{k \in \frac{n}{2} + \mathbb{Z}_+} z_2^{l+k-n+1} \sum_{j \in \mathbb{Z}_+} \left(\overset{\text{wt} a + q}{j} \right) \left(a_{j+l} b\right) *_{g,m,n}^k v + O_{g,k,m}''(V) \\ &= \sum_{k \in \frac{n}{2} + \mathbb{Z}_+} z_2^{l+k-n+1} \left(\operatorname{Res}_z \frac{(1 + z)^{\operatorname{wt} a + q}}{z^{-l}} Y(a, z) b\right) *_{g,m,n}^k v + O_{g,k,m}''(V) \\ &= \sum_{k \in \frac{n}{2} + \mathbb{Z}_+} z_2^{l+k-n+1} \operatorname{Res}_z \left(f_{-l}(a, b) *_{g,m,n}^k v \right) + O_{g,k,m}''(V) \qquad \text{(by Notation 6.1(3))} \\ &= \sum_{k \in \frac{n}{2} + \mathbb{Z}_+} z_2^{l+k-n+1} \sum_{j=0}^{k+1+q+l} (-1)^j \binom{l}{j} \sum_{i=0}^{k+1+q+l-j} (-1)^i \binom{-l+i+j-1}{i} \\ &\times \operatorname{Res}_z \left(f_{i+j-l}(a, b) *_{g,m,n}^k v \right) + O_{g,k,m}''(V) \qquad \text{(by Notation 6.1 (2)-(3) and Lemma 6.2)} \\ &= \sum_{k \in \frac{n}{2} + \mathbb{Z}_+} z_2^{l+k-n+1} \sum_{j=0}^{k+1+q+l} (-1)^j \binom{l}{j} \left(\left(a *_{g,n,k+1+q+l-j}^k b\right) *_{g,m,n}^k v \right) + O_{g,k,m}''(V) \\ \qquad \text{(by Lemma 6.3)} \end{aligned}$$

$$= \sum_{\substack{k \in \frac{\alpha}{T} + \mathbb{Z}_+ \\ k+1+l+q > 0}} z_2^{l+k-n+1} \sum_{j \in \mathbb{Z}_+} (-1)^j \binom{l}{j} a *_{g,m,k+1+q+l-j}^k \left(b *_{g,m,n}^{k+1+q+l-j} v \right) + O_{g,k,m}''(V)$$

(by (2.5) and Notation 6.1(1)-(2))

$$= \sum_{j \in \mathbb{Z}_{+}} \sum_{\substack{-n \leq i \in -s/T + \mathbb{Z} \\ -l+i+j \geq 1 + q - n}} \binom{l}{j} (-1)^{j} z_{2}^{i+j-q} a *_{g,m,n+i}^{-l+i+j-1-q+n} \left(b *_{g,m,n}^{n+i} v \right) \\ + O_{g,-l+i+j-1-q+n,m}''(V) \\ = \sum_{j \in \mathbb{Z}_{+}} \binom{l}{j} (-1)^{j} a_{\text{wt } a+q+l-j} \sum_{-n \leq i \in -s/T + \mathbb{Z}} z_{2}^{i+j-q} b_{\text{wt } b-1-i} (v + O_{g,n,m}''(V)) \\ = \sum_{j \in \mathbb{Z}_{+}} \binom{l}{j} (-1)^{j} a_{\text{wt } a+q+l-j} z_{2}^{\text{wt } b+j-q} Y_{M_{g}^{(m)}} \left(b, z_{2} \right) \left(v + O_{g,n,m}''(V) \right) \\ = \operatorname{Res}_{z_{0}} z_{0}^{l} \left(z_{0} + z_{2} \right)^{\text{wt } a+q} z_{2}^{\text{wt } b-q} Y_{M_{g}^{(m)}} \left(a, z_{0} + z_{2} \right) Y_{M_{g}^{(m)}} \left(b, z_{2} \right) \left(v + O_{g,n,m}''(V) \right),$$
 proving (3). \square

As an immediate consequence of Lemma 6.4 and Theorem 2.1 we have:

Proposition 6.5. For any $m \in (1/T)\mathbb{Z}_+$, $M_g^{(m)}$ is an admissible g-twisted V-module.

Recall from Section 2 that

$$L_{n,m}(V) = \operatorname{span} \{ (L(-1) + L(0) + m - n)u \mid u \in V \}.$$

From [4, Lemma 3.1] and the definition of $O'_{g,n,m}(V)$, we know:

Lemma 6.6. For any $m, n \in (1/T)\mathbb{Z}_+$,

$$\bigoplus_{s \not\equiv \bar{m} - \bar{n} \bmod T} V^s + L_{n,m}(V) \subseteq O'_{g,n,m}(V).$$

For any $u, v \in V$, it follows from [28] (see also [4]) that

$$Y(v,z)u \equiv (1+z)^{-\operatorname{wt} u - \operatorname{wt} v - m + n} Y\left(u, \frac{-z}{1+z}\right) v \bmod L_{n,m}(V).$$

Then the following result, in fact, was proved already in [4, Lemma 3.4 and Corollary 3.5].

Lemma 6.7. For $u \in V^r$ and $v \in V^s$, if $\bar{p} - \bar{n} \equiv r \mod T$, $\bar{m} - \bar{p} \equiv s \mod T$ and $m + n - p \ge 0$, then

$$u *_{a,m,p}^{n} v - v *_{a,m,m+n-p}^{n} u - \operatorname{Res}_{z}(1+z)^{\operatorname{wt} u - 1 + p - n} Y(u,z) v \in L_{n,m}(V).$$

In particular, taking p = m and v = 1 we have

$$u *_{q,m}^{n} \mathbf{1} - u \in L_{n,m}(V).$$

Theorem 6.8. For any $m, n \in (1/T)\mathbb{Z}_+$,

$$O_{g,n,m}(V) = \bigoplus_{s \not\equiv \bar{m} - \bar{n} \bmod T} V^s + L_{n,m}(V) + O''_{g,n,m}(V).$$

In particular, $O_{1,n,m}(V) = L_{n,m}(V) + O''_{1,n,m}(V)$.

Proof. By Proposition 6.5 and Theorem 3.5 (3), $V/O_{g,m,m}''(V) \subseteq \Omega_m(M_g^{(m)})$. Note by the definition of $O_{g,n,m}(V)$ that $O_{g,n,m}(V) = \bigoplus_{r=0}^{T-1} (O_{g,n,m}(V) \cap V^r)$. For any $u \in O_{g,n,m}(V) \cap V^r = \mathcal{O}_{g,n,m}(V) \cap V^r$ (see Theorem 5.2), then by the definition of $\mathcal{O}_{g,n,m}(V)$,

$$0 = o_{m-n}(u)(\mathbf{1} + O''_{g,m,m}(V)) = u *_{g,m}^{n} \mathbf{1} + O''_{g,n,m}(V),$$

i.e.,

$$u *_{g,m}^n \mathbf{1} \in O_{g,n,m}''(V).$$

If $\bar{m} - \bar{n} \equiv r \mod T$, then by Lemma 6.7

$$u = u - u *_{a,m}^{n} \mathbf{1} + u *_{a,m}^{n} \mathbf{1} \in L_{n,m}(V) + O''_{a,n,m}(V);$$

otherwise, $u \in \bigoplus_{s \not\equiv \bar{m} - \bar{n} \bmod T} V^s$. Thus by Lemma 6.6,

$$O_{g,n,m}(V) = \bigoplus_{s \not\equiv \bar{m} - \bar{n} \bmod T} V^s + L_{n,m}(V) + O''_{g,n,m}(V).$$

And when g = 1, it is clear that $O_{1,n,m}(V) = L_{n,m}(V) + O''_{1,n,m}(V)$. \square

7. Appendix

A detailed proof of Theorem 2.1 is given in this appendix.

Proposition 7.1. The twisted Jacobi identity (2.1) for $u \in V^r, v \in V^s$ and $w \in M$ is equivalent to: for any $w' \in M^* = \text{Hom}(M, \mathbb{C})$, there exist $l \in \mathbb{Z}_+, f(z_1, z_2) \in \mathbb{C}((z_1, z_2))$ and $g(z_0, z_2) \in \mathbb{C}((z_0, z_2))$ such that

$$\left\langle w', z_1^{\frac{r}{T}} z_2^{\frac{s}{T}} Y_M(u, z_1) Y_M(v, z_2) w \right\rangle = f(z_1, z_2) (z_1 - z_2)^{-l},$$
 (D1)

$$\left\langle w', z_1^{\frac{r}{T}} z_2^{\frac{s}{T}} Y_M(v, z_2) Y_M(u, z_1) w \right\rangle = f(z_1, z_2) (-z_2 + z_1)^{-l},$$
 (D2)

$$\left\langle w', (z_2 + z_0)^{\frac{r}{T}} z_2^{\frac{s}{T}} Y_M \left(Y(u, z_0) v, z_2 \right) w \right\rangle = g(z_0, z_2) (z_2 + z_0)^{-l},$$
 (D3)

$$\left\langle w', (z_0 + z_2)^{\frac{r}{T}} z_2^{\frac{s}{T}} Y_M(u, z_0 + z_2) Y_M(v, z_2) w \right\rangle = g(z_0, z_2) (z_0 + z_2)^{-l}, \quad (D4)$$

$$f(z_2 + z_0, z_2) z_0^{-l} = g(z_0, z_2) (z_2 + z_0)^{-l}$$
 (D5)

Proof. Assume that there exist $l \in \mathbb{Z}_+$, $f(z_1, z_2)$ and $g(z_0, z_2)$ such that (D1)-(D5) hold. Recall from [5,13] that

$$z_1^{-1} \left(\frac{z_2 + z_0}{z_1} \right)^{\gamma} \delta \left(\frac{z_2 + z_0}{z_1} \right) = z_2^{-1} \left(\frac{z_1 - z_0}{z_2} \right)^{-\gamma} \delta \left(\frac{z_1 - z_0}{z_2} \right)$$
(7.1)

for $\gamma \in \mathbb{C}$. Thus, (2.1) is equivalent to: for any $w' \in M^*$,

$$\begin{split} &z_{0}^{-1}\delta\left(\frac{z_{1}-z_{2}}{z_{0}}\right)\left\langle w',z_{1}^{\frac{r}{T}}z_{2}^{\frac{s}{T}}Y_{M}\left(u,z_{1}\right)Y_{M}\left(v,z_{2}\right)w\right\rangle \\ &-z_{0}^{-1}\delta\left(\frac{-z_{2}+z_{1}}{z_{0}}\right)\left\langle w',z_{1}^{\frac{r}{T}}z_{2}^{\frac{s}{T}}Y_{M}\left(v,z_{2}\right)Y_{M}\left(u,z_{1}\right)w\right\rangle \\ &=z_{1}^{-1}\delta\left(\frac{z_{2}+z_{0}}{z_{1}}\right)\left\langle w',\left(z_{2}+z_{0}\right)^{\frac{r}{T}}z_{2}^{\frac{s}{T}}Y_{M}\left(Y\left(u,z_{0}\right)v,z_{2}\right)w\right\rangle, \end{split}$$

which can be rewritten as

$$z_0^{-1}\delta\left(\frac{z_1-z_2}{z_0}\right)f\left(z_1,z_2\right)\left(z_1-z_2\right)^{-l}-z_0^{-1}\delta\left(\frac{-z_2+z_1}{z_0}\right)f\left(z_1,z_2\right)\left(-z_2+z_1\right)^{-l}$$

$$=z_1^{-1}\delta\left(\frac{z_2+z_0}{z_1}\right)f\left(z_2+z_0,z_2\right)z_0^{-l}$$

according to (D1)-(D3) and (D5). But this follows immediately from multiplying

$$z_0^{-1}\delta\left(\frac{z_1-z_2}{z_0}\right)-z_0^{-1}\delta\left(\frac{-z_2+z_1}{z_0}\right)=z_1^{-1}\delta\left(\frac{z_2+z_0}{z_1}\right)$$

by $f(z_1, z_2) z_0^{-l}$. Therefore, we get (2.1).

Conversely, assume that the commutator formula (2.3) and the weak associativity (2.2) hold. Moreover, we may choose l large enough such that $z^{l+\frac{r}{T}}Y_M(u,z)w$ involves only nonnegative integral powers of z. Then, it follows from (2.3) and (7.1) that

$$\left[Y_{M}\left(u,z_{1}\right),Y_{M}\left(v,z_{2}\right)\right]=\operatorname{Res}_{z_{0}}Y_{M}\left(z_{2}^{-1}\delta\left(\frac{z_{1}-z_{0}}{z_{2}}\right)Y\left(u,z_{0}\right)v,z_{2}\right)\left(\frac{z_{1}-z_{0}}{z_{2}}\right)^{-\frac{r}{T}}$$

$$\begin{split} &= \operatorname{Reg}_{z_{0}} Y_{M} \left(\left(z_{0}^{-1} \delta \left(\frac{z_{1} - z_{2}}{z_{0}} \right) - z_{0}^{-1} \delta \left(\frac{-z_{2} + z_{1}}{z_{0}} \right) \right) Y \left(u, z_{0} \right) v, z_{2} \right) \left(\frac{z_{1} - z_{0}}{z_{2}} \right)^{-\frac{r}{T}} \\ &= \operatorname{Res}_{z_{0}} Y_{M} \left(z_{0}^{-1} \delta \left(\frac{z_{1} - z_{2}}{z_{0}} \right) Y \left(u, z_{1} - z_{2} \right) v \right. \\ &\qquad \left. - z_{0}^{-1} \delta \left(\frac{-z_{2} + z_{1}}{z_{0}} \right) Y \left(u, -z_{2} + z_{1} \right) v, z_{2} \right) \left(\frac{z_{1} - z_{0}}{z_{2}} \right)^{-\frac{r}{T}} \\ &= z_{2}^{\frac{r}{T}} z_{1}^{-\frac{r}{T}} \sum_{n \geq 0} \operatorname{Res}_{z_{0}} Y_{M} \left(z_{0}^{-1} \delta \left(\frac{z_{1} - z_{2}}{z_{0}} \right) Y \left(u, z_{1} - z_{2} \right) v \right. \\ &\qquad \left. - z_{0}^{-1} \delta \left(\frac{-z_{2} + z_{1}}{z_{0}} \right) Y \left(u, -z_{2} + z_{1} \right) v, z_{2} \right) \left(\frac{-r/T}{n} \right) \left(\frac{-z_{0}}{z_{1}} \right)^{n} \\ &= z_{2}^{\frac{r}{T}} z_{1}^{-\frac{r}{T}} \sum_{n = 0}^{N} (-1)^{n} \binom{-r/T}{n} z_{1}^{-n} Y_{M} \left(\left(Y \left(u, z_{1} - z_{2} \right) \left(z_{1} - z_{2} \right)^{n} \right. \\ &\qquad \left. - Y \left(u, -z_{2} + z_{1} \right) \left(-z_{2} + z_{1} \right)^{n} \right) v, z_{2} \right), \end{split}$$

where N is a nonnegative integer such that $z^{N+1}Y(u,z)v \in V[[z]]$. Thus, there exists $l \in \mathbb{Z}_+$ such that

$$(z_1 - z_2)^l [Y_M(u, z_1), Y_M(v, z_2)] = 0.$$

Then for any $w \in M$,

$$(z_{1} - z_{2})^{l} \left\langle w', z_{1}^{\frac{r}{T}} z_{2}^{\frac{s}{T}} Y_{M}(u, z_{1}) Y_{M}(v, z_{2}) w \right\rangle$$

$$= (-z_{2} + z_{1})^{l} \left\langle w', z_{1}^{\frac{r}{T}} z_{2}^{\frac{s}{T}} Y_{M}(v, z_{2}) Y_{M}(u, z_{1}) w \right\rangle,$$

which is denoted by $f(z_1, z_2)$. Note that both sides of the above formula involve only finitely many negative powers of z_2 and z_1 . Thus, $f(z_1, z_2) \in \mathbb{C}((z_1, z_2))$, proving (D1) and (D2).

Set $g(z_0, z_2) = (z_0 + z_2)^l f(z_0 + z_2, z_2) z_0^{-l}$. Since $z^{l + \frac{r}{T}} Y_M(u, z) w \in M[[z]], z_1^l f(z_1, z_2) \in \mathbb{C}[[z_1, z_2, z_2^{-1}]]$, we obtain $g(z_0, z_2) \in \mathbb{C}((z_0, z_2))$. Then,

$$\begin{split} & \left\langle w', \left(z_{0}+z_{2}\right)^{\frac{r}{T}}z_{2}^{\frac{s}{T}}Y_{M}\left(u,z_{0}+z_{2}\right)Y_{M}\left(v,z_{2}\right)w\right\rangle \\ & = \left\langle w', z_{1}^{\frac{r}{T}}z_{2}^{\frac{s}{T}}Y_{M}\left(u,z_{1}\right)Y_{M}\left(v,z_{2}\right)w\right\rangle \Big|_{z_{1}=z_{0}+z_{2}} \\ & = f\left(z_{1},z_{2}\right)\left(z_{1}-z_{2}\right)^{-l}\Big|_{z_{1}=z_{0}+z_{2}} = f\left(z_{0}+z_{2},z_{2}\right)z_{0}^{-l} \\ & = \left(z_{0}+z_{2}\right)^{l}f\left(z_{0}+z_{2},z_{2}\right)z_{0}^{-l}\left(z_{0}+z_{2}\right)^{-l} = g\left(z_{0},z_{2}\right)\left(z_{0}+z_{2}\right)^{-l}, \end{split}$$

that is, (D4). Now by (2.2),

$$(z_0 + z_2)^l \left\langle w', (z_0 + z_2)^{\frac{r}{T}} z_2^{\frac{s}{T}} Y_M (u, z_0 + z_2) Y_M (v, z_2) w \right\rangle$$

$$= (z_2 + z_0)^l \left\langle w', (z_2 + z_0)^{\frac{r}{T}} z_2^{\frac{s}{T}} Y_M (Y (u, z_0) v, z_2) w \right\rangle = g (z_0, z_2) ,$$

from which one can deduce (D3) and (D5). \Box

Taking v = 1 in (2.2) we have

$$(z_2 + z_0)^{k + \frac{r}{T}} Y_M \left(e^{z_0 D} u, z_2 \right) w = (z_0 + z_2)^{k + \frac{r}{T}} Y_M \left(u, z_0 + z_2 \right) w, \tag{7.2}$$

where D is a linear operator on V defined by $D(v) = v_{-2}\mathbf{1}$ for $v \in V$. Note that we can choose l large enough such that $z^{l+\frac{r}{T}}Y_M(u,z)w \in M[[z]]$. It follows that (7.2) can also be written as

$$(z_2 + z_0)^{l + \frac{r}{T}} Y_M (e^{z_0 D} u, z_2) w = (z_2 + z_0)^{l + \frac{r}{T}} Y_M (u, z_2 + z_0) w.$$

Multiplying both sides by $(z_2 + z_0)^{-l - \frac{r}{T}}$ gives

$$Y_M \left(e^{z_0 D} u, z_2 \right) w = Y_M \left(u, z_2 + z_0 \right) w. \tag{7.3}$$

Now we are ready to present the proof of Theorem 2.1.

Proof. By Proposition 7.1, it is sufficient to deduce from the weak associativity that there exist $f(z_1, z_2) \in \mathbb{C}((z_1, z_2))$ and $g(z_0, z_2) \in \mathbb{C}((z_0, z_2))$ such that (D1)-(D5) hold. By (2.2) we have

$$(z_0 + z_2)^l \left\langle w', (z_0 + z_2)^{\frac{r}{T}} z_2^{\frac{s}{T}} Y_M (u, z_0 + z_2) Y_M (v, z_2) w \right\rangle$$

$$= (z_2 + z_0)^l \left\langle w', (z_2 + z_0)^{\frac{r}{T}} z_2^{\frac{s}{T}} Y_M (Y (u, z_0) v, z_2) w \right\rangle,$$

which is denoted by $g(z_0, z_2)$. Clearly, $g(z_0, z_2) \in \mathbb{C}((z_0, z_2))$. Then it is easy to see that (D3) and (D4) hold. Choose l to be sufficiently large such that $z_0^l g(z_0, z_2) \in \mathbb{C}[[z_0, z_2, z_2^{-1}]]$. Set $f(z_1, z_2) = (z_1 - z_2)^l g(z_1 - z_2, z_2) z_1^{-l}$, which lies in $\mathbb{C}((z_1, z_2))$. Then, $f(z_2 + z_0, z_2) z_0^{-l} = g(z_0, z_2) (z_2 + z_0)^{-l}$, proving (D5); and

$$\begin{split} & \left\langle w', z_{1}^{\frac{r}{T}} z_{2}^{\frac{s}{T}} Y_{M}\left(u, z_{1}\right) Y_{M}\left(v, z_{2}\right) w \right\rangle \\ & = \left\langle w', \left(z_{0} + z_{2}\right)^{\frac{r}{T}} z_{2}^{\frac{s}{T}} Y_{M}\left(u, z_{0} + z_{2}\right) Y_{M}\left(v, z_{2}\right) w \right\rangle \Big|_{z_{0} = z_{1} - z_{2}} \\ & = g\left(z_{0}, z_{2}\right) \left(z_{0} + z_{2}\right)^{-l} \Big|_{z_{0} = z_{1} - z_{2}} = g\left(z_{1} - z_{2}, z_{2}\right) z_{1}^{-l} = f(z_{1}, z_{2})(z_{1} - z_{2})^{-l}, \end{split}$$

proving (D1).

By (D1), there exists $F(z_1, z_2) \in \mathbb{C}((z_1, z_2))$ such that $\langle w', z_1^{\frac{r}{T}} z_2^{\frac{s}{T}} Y_M(v, z_2) Y_M(u, z_1) w \rangle = F(z_1, z_2) (-z_2 + z_1)^{-l}$ and $z_2^l F(z_1, z_2) \in \mathbb{C}[[z_1, z_1^{-1}, z_2]]$. Then,

$$\begin{split} & \left\langle w', z_{1}^{\frac{r}{T}} \left(-z_{0}+z_{1}\right)^{\frac{s}{T}} Y_{M} \left(v,-z_{0}+z_{1}\right) Y_{M} \left(u,z_{1}\right) w \right\rangle \\ & = \left\langle w', z_{1}^{\frac{r}{T}} z_{2}^{\frac{s}{T}} Y_{M} \left(v,z_{2}\right) Y_{M} \left(u,z_{1}\right) w \right\rangle \Big|_{z_{2}=-z_{0}+z_{1}} \\ & = F \left(z_{1},-z_{0}+z_{1}\right) z_{0}^{-l}. \end{split}$$

Thus,

$$\left\langle w', z_{1}^{\frac{r}{T}}\left(z_{1}-z_{0}\right)^{\frac{s}{T}}Y_{M}\left(Y\left(v,-z_{0}\right)u,z_{1}\right)w\right\rangle = \left(-z_{0}+z_{1}\right)^{l}F\left(z_{1},-z_{0}+z_{1}\right)z_{0}^{-l}\left(z_{1}-z_{0}\right)^{-l}$$

by the weak associativity. Then,

$$f(z_{2} + z_{0}, z_{2}) z_{0}^{-l} = g(z_{0}, z_{2}) (z_{2} + z_{0})^{-l}$$

$$= \left\langle w', (z_{2} + z_{0})^{\frac{r}{T}} z_{2}^{\frac{s}{T}} Y_{M} (Y(u, z_{0}) v, z_{2}) w \right\rangle \qquad \text{(by (D3))}$$

$$= \left\langle w', (z_{2} + z_{0})^{\frac{r}{T}} z_{2}^{\frac{s}{T}} Y_{M} (e^{z_{0}D} Y(v, -z_{0}) u, z_{2}) w \right\rangle$$

$$= \left\langle w', z_{2}^{\frac{s}{T}} (z_{2} + z_{0})^{\frac{r}{T}} Y_{M} (Y(v, -z_{0}) u, z_{2} + z_{0}) w \right\rangle \qquad \text{(by (7.3))}$$

$$= \left\langle w', z_{1}^{\frac{r}{T}} (z_{1} - z_{0})^{\frac{s}{T}} Y_{M} (Y(v, -z_{0}) u, z_{1}) w \right\rangle \Big|_{z_{1} = z_{2} + z_{0}}$$

$$= ((-z_{0} + z_{1})^{l} F(z_{1}, -z_{0} + z_{1}) z_{0}^{-l}) (z_{1} - z_{0})^{-l} \Big|_{z_{1} = z_{2} + z_{0}}$$

$$= F(z_{2} + z_{0}, z_{2}) z_{0}^{-l}.$$

Thus, $F(z_2 + z_0, z_2) = f(z_2 + z_0, z_2)$ and then $F(z_1, z_2) = f(z_1, z_2)$, that is, (D2). This completes the proof. \square

Acknowledgment

J. Han is supported by the National Natural Science Foundation of China (No. 12271406). Y. Xiao is supported by the National Natural Science Foundation of China (No. 12401034) and the Natural Science Foundation of Shandong Province (No. ZR2023QA066).

Data availability

No data was used for the research described in the article.

References

- R. Borcherds, Vertex algebras, Kac-Moody algebras, and the monster, Proc. Natl. Acad. Sci. USA 83 (1986) 3068–3071.
- C. Dong, J. Han, Some finite properties for vertex operator superalgebras, Pac. J. Math. 258 (2012) 269–290.
- [3] C. Dong, C. Jiang, Bimodules associated to vertex operator algebras, Math. Z. 259 (2008) 799–826.
- [4] C. Dong, C. Jiang, Bimodules and g-rationality of vertex operator algebras, Trans. Åm. Math. Soc. 360 (2008) 4235–4262.
- [5] C. Dong, J. Lepowsky, Generalized Vertex Algebras and Relative Vertex Operators, Progr. Math., vol. 112, Birkhäuser Boston, Inc., Boston, MA, 1993.
- [6] C. Dong, H. Li, G. Mason, Vertex operator algebras and associative algebras, J. Algebra 206 (1998) 67–96.
- [7] C. Dong, H. Li, G. Mason, Twisted representations of vertex operator algebras, Math. Ann. 310 (1998) 571–600.
- [8] C. Dong, H. Li, G. Mason, Twisted representations of vertex operator algebras and associative algebras, Int. Math. Res. Not. 8 (1998) 389–397.
- [9] C. Dong, L. Ren, Representations of vertex operator algebras and bimodules, J. Algebra 384 (2013) 212–226.
- [10] C. Dong, C. Yang, G-twisted associative algebras of vertex operator superalgebras, J. Algebra 606 (2022) 323–340.
- [11] C. Dong, Z. Zhao, Twisted representations of vertex operator superalgebras, Commun. Contemp. Math. 8 (2006) 101–122.
- [12] I. Frenkel, Y. Huang, J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, Mem. Am. Math. Soc. 104 (1993) 1–64.
- [13] I. Frenkel, J. Lepowsky, A. Meurman, Vertex Operator Algebras and the Monster, Pure Appl. Math., vol. 134, Academic Press, Boston, 1988.
- [14] I. Frenkel, Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, Duke Math. J. 66 (1992) 123–168.
- [15] J. Han, Bimodules and universal enveloping algebras associated to VOAs, Isr. J. Math. 247 (2022) 905–922.
- [16] J. Han, Y. Xiao, Associative algebras and universal enveloping algebras associated to VOAs, J. Algebra 564 (2020) 489–498.
- [17] X. He, Higher level Zhu algebras are subquotients of universal enveloping algebras, J. Algebra 491 (2017) 265–279.
- [18] Y. Huang, J. Yang, Logarithmic intertwining operators and associative algebras, J. Pure Appl. Algebra 216 (2012) 1467–1492.
- [19] Y. Huang, J. Yang, Associative algebras for (logarithmic) twisted modules for a vertex operator algebra, Trans. Am. Math. Soc. 371 (2019) 3747–3786.
- [20] V. Kac, W. Wang, Vertex operator superalgebras and representations, Contemp. Math. 175 (1994) 161–191.
- [21] J. Lepowsky, H. Li, Introduction to Vertex Operator Algebras and Their Representation, Progr. Math., vol. 227, Birkhäuser Boston, Inc., Boston, MA, 2004.
- [22] H. Li, Determining fusion rules by A(V)-modules and bimodules, J. Algebra 212 (1999) 515–556.
- [23] H. Li, Regular representations and $A_n(V)$ - $A_m(V)$ bimodules, J. Pure Appl. Algebra 227 (2023) 25pp.
- [24] M. Miyamoto, K. Tanabe, Uniform product of $A_{g,n}(V)$ for an orbifold model V and G-twisted Zhu algebra, J. Algebra 274 (2004) 80–96.
- [25] C. Yang, M. Yang, Associative algebras of Z-graded vertex operator superalgebras and their applications, J. Algebra 633 (2023) 20−42.
- [26] J. Yang, On associative algebras, modules and twisted modules for vertex operator algebras, J. Algebra 440 (2015) 354–378.
- [27] Y. Zhu, Bimodules over twisted Zhu algebras and a construction of tensor product of twisted modules for vertex operator algebras, preprint, arXiv:2409.08995 [math. QA], 2024.
- [28] Y. Zhu, Modular invariance of characters of vertex operator algebras, J. Am. Math. Soc. 9 (1996) 237–302.